

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/15, 31/155, C07C 243/20, 247/14, 279/12, 281/16		A1	(11) International Publication Number: WO 97/02027
			(43) International Publication Date: 23 January 1997 (23.01.97)
(21) International Application Number: PCT/US96/11318		Aharoni Street, 76281 Rehovot (IL). MILMAN, Isaac [IL/IL]; 36/37 Hakeren, 90610 Maale Adumim (IL). FINKELSTEIN, Nina [IL/IL]; 119/16 Neve Yaacov, 97350 Jerusalem (IL). (74) Agent: WHITE, John, P.; Cooper & Dunham L.L.P., 1185 Avenue of the Americas, New York, NY 10036 (US). (81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 5 July 1996 (05.07.96)			
(30) Priority Data: 08/498,095 5 July 1995 (05.07.95) US			
(60) Parent Application or Grant (63) Related by Continuation US 08/498,095 (CIP) Filed on 5 July 1995 (05.07.95)			
(71) Applicant (for all designated States except US): TEVA PHARMACEUTICAL INDUSTRIES, LTD. [IL/IL]; Science Based Industries Campus, Har Hotzvim, P.O. Box 1142, 91010 Jerusalem (IL).		Published With international search report.	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): GILAD, Gad [IL/IL]; Rachel Street, 53482 Givatayim (IL). GILAD, Varda [IL/IL]; Rachel Street, 53482 Givatayim (IL). STERLING, Jeffrey [US/IL]; 156/1 Ramot, 97275 Jerusalem (IL). HERZIG, Yaacov [IL/IL]; 7 Hashiriyon Street, 43266 Raanana (IL). LERNER, David [US/IL]; 27/4 Yigal Street, 97241 Jerusalem (IL). VEINBERG, Alex [IL/IL]; 5/16			
(54) Title: POLYAMINE DERIVATIVES OF 1-AMINOINDAN			
(57) Abstract			
<p>A compound of general formula (I), wherein n is 1 or 2; R₁ and R₂ are each independently hydrogen, hydroxy, substituted or unsubstituted C₁₋₄ alkyl, substituted or unsubstituted C₁₋₄ alkoxy or halogeno; R₅ is hydrogen, substituted or unsubstituted C₁₋₄ alkyl, amino or NR₆R₇; R₃, R₄, R₆ and R₇ are each independently hydrogen or (CH₂)_m-[NH(CH₂)_x]_y-NHR₁₁ wherein m is from 0-5, y is from 0-5 and each x is independently from 1-5 and R₁₁ is hydrogen or C(NH)NH₂; provided that when R₃ and R₄ are both hydrogen, R₅ is amino or NR₆R₇; pharmaceutically acceptable salts and optically active isomers thereof.</p>			
		<p style="text-align: right;">(I)</p>	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

POLYAMINE DERIVATIVES OF 1-AMINOINDAN

5 Throughout this application, various references are referred to. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains.

10

Background of the Invention

The present invention relates to novel polyamines, compositions containing them and their use in the treatment of neurotrauma such as stroke.

15

Polyamines as a class of compounds have been reported to possess a wide range of activities both within and external to the central nervous system. Such activities include modulation of postsynaptic receptors, such as nicotinic, glutamic and N-methyl-D-aspartate receptors, modulation of calcium regulatory processes as well as antiplatelet, antiinflammatory and anticoagulant activity.

25


Various polyamine containing molecules have been synthesized and screened in a variety of animal models of disorders of the central nervous system. Many of these compounds have been based on toxins extracted from spiders such as A. aperta and Argiope labata and the wasp P. triangulum. These compounds comprise a long poly(aminoalkane) chain linked to an aromatic or heterocyclic group through a carbonyl group.

30


35 Guanidine derivatives of polyamines, e.g. SFTX-33 have been reported as voltage-sensitive calcium channel blockers and EAA antagonists capable of binding at the PCP site on the NMDA receptor. U.S. Patent No. 4,906,779 (Oregon 1990). Spergualine and

-2-

deoxypergualine has been reported (R.J. Bergeron, J.S. McManis, J. Org. Chem., 52, 1700 (1987); Drug Fut., 16, 165 (1991)) to possess immunosuppressant activity and beneficial effect in the treatment of Multiple Sclerosis.



sFTX-33



spergualine (R=OH)
deoxyspergualine (R=H)

Gilad G et al. (Life Sci. (1989) 44 1963-1969 and Exp. Neurol. (1991) 111 349-355) have described activity of the polyamines spermine, spermidine and putrescine in protecting against ischemia induced nerve cell death in gerbils.

Disadvantages associated with using the simple polyamines described in Gilad G et al. relate to transport through biological membranes and various toxic side effects that would hinder these compounds from proceeding to clinical use.


Accordingly, the present invention provides derivatives of spermine, spermidine, and putrescine and guanidine

-3-

derivatives thereof that retain the activity of the polyamine but display fewer undesirable side effects.

Summary of the Invention

This invention provides compounds of general formula I



(I)

wherein n is 1 or 2;

R₁ and R₂ are each independently hydrogen, hydroxy, substituted or unsubstituted C₁₋₄ alkyl, substituted or unsubstituted C₁₋₄ alkoxy or halogeno;

R₅ is hydrogen, substituted or unsubstituted C₁₋₄ alkyl, amino or NR₆R₇;

R₃, R₄, R₆ and R₇ are each independently hydrogen or (CH₂)_m-[NH(CH₂)_x]_y-NHR₁₁ wherein m is from 0-5, y is from 0-5 and each x is independently from 1-5, R₁₁ is hydrogen or C(NH)NH₂;

provided that when R₃ and R₄ are both hydrogen, R₅ is amino or NR₆R₇; pharmaceutically acceptable salts and optically active isomers thereof.

Description of Figures


- Figure 1A: Scheme I - Schematic of chemical synthesis Examples 1-5.
- Figure 1B: Scheme I continued.
- Figure 2: Scheme II - Schematic of chemical synthesis Example 6.
- Figure 3: Scheme III - Schematic of chemical synthesis of Example 7.
- Figure 4: Schematic of chemical synthesis of Example 8.

Detailed Description of the Invention

This invention provides compounds of general formula I

5

10



(I)

wherein n is 1 or 2;

15

R_1 and R_2 are each independently hydrogen, hydroxy, substituted or unsubstituted C_{1-4} alkyl, substituted or unsubstituted C_{1-4} alkoxy or halogeno;

20

R_5 is hydrogen, substituted or unsubstituted C_{1-4} alkyl, amino or NR_6R_7 ;

R_3 , R_4 , R_6 and R_7 are each independently hydrogen or $(CH_2)_m-[NH(CH_2)_x]_y-NHR_{11}$ wherein m is from 0-5, y is from 0-5 and each x is independently from 1-5, R_{11} is hydrogen or $C(NH)NH_2$;

provided that when R_3 and R_4 are both hydrogen, R_5 is amino or NR_6R_7 ; pharmaceutically acceptable salts and optically active isomers thereof.

25

30

Within general formula I it is generally preferred that n is 1, that is the compounds are derivatives of 1-aminoindan. It is further preferred that R_1 and R_2 are each independently hydrogen, hydroxy or halogeno, most preferably R_1 and R_2 are both hydrogen. In a more specific embodiment R_1 , R_2 and R_5 are hydrogen.

35

In one embodiment of the invention one of R_3 and R_4 is hydrogen and the other is of the formula $(CH_2)_m-[NH(CH_2)_x]_y-NHR_{11}$. In such an embodiment it is preferable that R_{11} is hydrogen, m is 3 or 4, y is 0, 1 or 2 and x where appropriate is 3 or 4. When y is more than 1 it is

-7-

preferable that each x is 3 or 4.

Alternatively, one of R_3 and R_4 are hydrogen and the other is of the formula $(CH_2)_m-[NH(CH_2)_x]_y-NH-C(NH)NH_2$,
5 i.e. R_{11} is $C(NH)NH_2$, m, y and x being as defined above.

In an alternative embodiment both R_3 and R_4 are of the formula $(CH_2)_m-[NH(CH_2)_x]_y-NHR_{11}$. In this alternative embodiment it is preferable that m is 3 or 4 and y is 0,
10 1 or 2 more preferably 0. In a specific embodiment, R_{11} is H.

In a further alternative embodiment of the invention, both R_3 and R_4 are hydrogen and R_5 is amino or NR_6R_7 ,
15 wherein R_6 and R_7 are each independently hydrogen or of the formula $(CH_2)_m-[NH(CH_2)_x]_y-NHR_{11}$ with m, x, y and R_{11} taking the preferences described above. Preferably R_5 is amino or $N(CH_2)_m-[NH(CH_2)_x]_y-NHR_{11}$ with m being 3 or 4, y being 0 and R_{11} , hydrogen.

20 The compounds of formula I are aminoindan or aminotetralin analogs of spermine, spermidine, and putrescine and guanidine derivatives thereof and isomers of these polyamines. Many of the compounds of general
25 formula I are optically active. The present invention further relates to the purified optically active isomers.

In such embodiments when R_5 is not hydrogen, the R_5 group
30 may exist in either the cis or trans configuration with respect to the group NR_3R_4 . In such circumstances it is preferable that R_5 and the group NR_3R_4 are in the cis configuration.

35 In all the above embodiments, the ring carbon to which the NR_3R_4 group is bound may be in the "R" or "S" configuration, preferably it is in the "R"

configuration.

Specific embodiments of the compound of general formula I include the following: N-(3-aminopropyl)-1-aminoindan;

- 5 N-(4-aminopropyl)-1-aminoindan;
N,N-di(4-amino-n-butyl)-1-aminoindan;
(R)-N,N-di(4-amino-n-butyl)-1-aminoindan;
(S)-N,N-di(4-amino-n-butyl)-1-aminoindan;
N-(4-amino-n-butyl)-N-(3-amino-n-propyl)-1-aminoindan;
10 N,N-di(4-amino-n-butyl)-1-amino-1,2,3,4-tetrahydronaphthalene;
N-[N-(4-aminobutyl)-3-aminopropyl]-1-aminoindan;
N-[N-(3-aminopropyl)-4-aminobutyl]-1-aminoindan;
N-{[N-(3-aminopropyl)-4-aminobutyl]}-N'-3-aminopropyl}-
15 1-aminoindan;
N-{[N-(4-aminobutyl)-3-aminopropyl]}-N'-4-aminobutyl}-1-aminoindan;
cis N,N'-diacetyl-1,3-diaminoindan;
trans N,N'-diacetyl-1,3-diaminoindan;
20 cis 1,3-diaminoindan;
trans 1,3-diaminoindan;
N-(4-aminobutyl)-1,3-diaminoindan;
N,N'-di(4-aminobutyl)-1,3-diaminoindan;
N-(3-aminopropyl)-N-[N-(3-aminopropyl)-4-aminobutyl]-1-
25 aminoindan;
N,N-di(4-guanidinobutyl)-1-aminoindan;
N-(4-guanidinobutyl)-1-aminoindan, and pharmaceutically acceptable salts thereof.

- 30 The present invention also provides a pharmaceutical composition comprising a therapeutically effective amount of at least one of the above-listed compounds and a pharmaceutically acceptable carrier.

- 35 This invention provides for the pharmaceutical composition wherein the pharmaceutically acceptable carrier is a solid and the pharmaceutical composition is

-9-

a tablet. In an embodiment the therapeutically effective amount is from about 1 mg to about 1000 mg. In a more specific embodiment the therapeutically effective amount is from about 10 mg to about 100 mg.

5

In another embodiment of the pharmaceutical composition the pharmaceutically acceptable carrier is a liquid and the pharmaceutical composition is an injectable solution. In a specific embodiment the therapeutically effective amount is from about 1 mg/ml to about 1000 mg/ml. In a more specific embodiment the therapeutically effective amount is from about 10 mg/ml to about 100 mg/ml.

10

15 In an embodiment of the pharmaceutical composition the carrier is a gel and the pharmaceutical composition is a suppository.

This invention further provides a method for treating acute neurological traumatic disorder or neurotrauma in a subject comprising administering to the subject a therapeutically effective amount of a compound of the formula I or pharmaceutically acceptable salt thereof, effective to treat the acute neurological traumatic disorder or neurotrauma on the subject.

20

25

The subject can be any vertebrate, but is preferably a mammal, for example a cow, goat, sheep, horse, dog, monkey, ape, rabbit, rat or mouse. Most preferably the mammalian subject is a human.

30

The subject invention further provides a method of treating a subject afflicted with a neurodegenerative disease which comprises administering to the subject an amount of a compound of general formula I or the pharmaceutically acceptable salt thereof of the subject invention effective to treat the neurodegenerative

35

disease in the subject.

5 The subject invention further provides a method of treating a subject afflicted with a neurotoxic injury which comprises administering to the subject an amount of a compound of general formula I or the pharmaceutically acceptable salt thereof of the subject invention effective to treat the neurotoxic injury in the subject.

10

The subject invention further provides a method of treating a subject afflicted with brain ischemia which comprises administering to the subject an amount of a compound of general formula I or the pharmaceutically acceptable salt thereof of the subject invention effective to treat brain ischemia in the subject.

15

20 The subject invention further provides a method of treating a subject afflicted with a head trauma injury which comprises administering to the subject an amount of a compound of general formula I or the pharmaceutically acceptable salt thereof of the subject invention effective to treat the head trauma injury in the subject.

25

The subject invention further provides a method of treating a subject afflicted with a spinal trauma injury which comprises administering to the subject an amount of a compound of general formula I or the pharmaceutically acceptable salt thereof of the subject invention effective to treat the spinal trauma injury in the subject.

30

35 The subject invention further provides a method of preventing nerve damage in a subject which comprises administering to the subject an amount of general formula I or the pharmaceutically acceptable salt

-11-

thereof of the subject invention effective to prevent nerve damage in the subject.

5 The terms acute neurological traumatic disorder or neurotrauma are used herein to refer to, and include clinical indications such as strokes, acute cranial injury.

10 The compounds of the present invention are also of use as immunosuppressants, as anti-neoplastic agents, anti-psoratic agents and in the treatment of epilepsy, Alzheimer's Disease, multiple sclerosis, as insecticides and pesticides against endo- and ectoparasites.

15 The compound of general formula I, its salts and optically active isomers can be administered using techniques well known in the art. For example, the compound may be administered orally, rectally, parenterally, intravenously, or intramuscularly.

20 To aid the discussion of the preparation of the compounds of the present invention it is convenient to classify them as follows:

25 1) Diamines (e.g. derivatives of putrescine or analogues with a different number of methylene groups in the alkyl chain;

30 2) Triamines (e.g. derivatives of spermidine or analogues with differing numbers of methylene groups in the alkyl chain;


35 3) Tetramines (e.g. derivatives of spermine or analogues with differing numbers of methylene groups in the alkyl chain;

4) Compounds with five or more amino groups which are

-12-

not direct derivatives of natural polyamines.

With reference to the schemes diagrammed in Figures 1 and 2, the compounds of the present invention may be prepared from the parent 1-aminoindan or 1-aminotetralin by reaction with a compound of the formula A or B.



wherein m is defined as previously and X is a suitable leaving group (such as halide or sulfonate ester) to enable N-alkylation by A or B; and one or both of R_8 and R_9 are suitable amino protecting groups such as t-butyloxycarboxy (BOC) or together are phthaloyl. Alternatively the terminal nitrile group of compound B fulfills the same role as an amine protecting group.

The reaction between the 1-aminoindan or 1-aminotetralin can be controlled to give predominantly either diamines or symmetrical triamines or a mixture of the two which may be separated into pure compounds by methods known to those skilled in the art (chromatography, crystallization, etc.). The desired diamines or symmetrical triamines may be isolated by removing the protecting groups under the appropriate conditions. For example if R_8 in IA is BOC, treatment with hydrogen chloride in ether. If R_8R_9 in IA is phthaloyl, treatment with hydrazine. If the protecting group is nitrile in IB it is reduced with lithium aluminum hydride (LAH). In general the diamine is isolated as the salt of a suitable acid (e.g. hydrochloric, sulfuric, methansulfonic, etc.).

Unsymmetrical triamines may be prepared by reacting a

-13-

second alternative compound of the formula A or B with the partially protected diamine (I). The terminal protecting groups may then be removed according to methods described immediately above.

5

Other unsymmetrical triamines may be prepared by reacting a second alternative compound of the formula A or B and the suitably protected diamine (III) which is prepared from partially protected diamine (I) by
10 introducing a second protecting group on the secondary amine to give fully protected diamine (II). The second protecting group (R_{10}) is chosen so that the terminal primary amine protecting group (R_8R_9 or nitrile) may be selectively converted to a free amine
15 in its presence. For example, if R_8R_9 is phthaloyl, R_{10} may be BOC. In the case where R_8R_9 of the fully protected diamine (IIA) is phthaloyl, it may be selectively removed by treatment with hydrazine to give a compound of formula (III). In the case where
20 the fully protected diamine (IIB) has a nitrile group, it may be selectively reduced with LAH likewise to give a compound of formula (III). In either case, the resulting compound of formula (III) may be reacted with a compound of formula A or B as described above
25 to give a partially protected triamine of formula (IV). The desired triamine may be isolated as a suitable salt by removal of the protecting groups under the appropriate conditions as described above and subsequent reaction with a suitable acid.

30

Alternatively, the partially protected triamine (IV) may be converted to a tetramine by repeating the process described above (full protection to (V) and selective deprotection to (VI). The resulting
35 partially protected tetramine (VII) may be fully deprotected to isolate the tetramine or the protection/partial deprotection may be repeated to a

pentamine, or higher homologue.



In a further embodiment of the invention the indan or tetralin skeleton forms an integral part of the polyamine structure. In this case the diamine
5 compound may be 1,3-diaminoindan (or 1,3-diaminotetralin) or 1,4-diaminotetralin. Each analogue may be isolated as the cis or trans isomer. They may be prepared from the corresponding diketone
10 by reaction with hydroxylamine hydrochloride followed by reduction (preferably using sodium in alcohol (preferably ethanol or isopropanol) or catalytic hydrogenation, preferably with palladium on carbon. The resulting cis/trans mixture of diamines may be
15 resolved by any convenient means, preferably by conversion to the diacetyl amide and resolution by crystallization and/or chromatography. The isomerically pure diacetates may be hydrolyzed to the desired diamines under acidic conditions (preferably
20 aqueous hydrochloric acid).

In analogy to the procedures described above, these diamines may be converted to triamines and tetramines by reaction with compounds of the formula A or B
25 (preferably B). In general, such a reaction results in a mixture of triamine (XI) and tetramine (XII) which may be separated by any convenient means, preferably by chromatography after converting them to the fully protected analogues (XIII) and (XIV), R_{10}
30 preferably is BOC. After resolution, the protecting groups are removed from the resulting pure compounds (XIII) and (XIV) preferably using LAH for the nitrile and hydrogen chloride in ether for the BOC. The desired compounds (XV) are thus isolated as a suitable
35 salt (hydrochloride, sulfuric acid, methanesulfonic acid etc) as relatively pure cis or trans isomers.

-15-

In a further embodiment of the invention, tetramines of the formula XVI can alternatively be prepared directly from the indan or tetralin diketone by reaction with a mono-protected diamine of formula C or

5 D



10 where m, R₈ and R₉ are all as defined previously, to give the corresponding imine (XVII) which may be reduced (preferably with sodium borohydride or by catalytic hydrogenation) followed by deprotection under conditions appropriate for the protecting group, preferably acid hydrolysis for the case where R₈ is
15 BOC. In this case XVI is isolated as a mixture of cis and trans isomers.

Unsymmetrical tetramines, e.g. compound 3069, may be
20 prepared from the partially protected triamine (XXII) by introducing an appropriately protected aminoalkyl group, followed by removal of the protecting groups.

The triamine XXII may be prepared from 1-aminoindan by
25 stepwise alkylations, starting with a protected aminoalkyl group, followed by a cyanoalkyl moiety. The nitrile group may subsequently be reduced with LAH and protected to give XXII.

30 The guanidine derivatives (compounds 3042, 3043) may be prepared by reacting the corresponding diamines or triamines with a guanylating agent e.g. N,N-di-tert-butoxycarbonylthiourea, (Iwanowicz E et al. Synthetic Communications (1993) 23, 1443), in a polar aprotic
35 medium e.g. DMF, in the presence of an organic base e.g. triethylamine, optionally in the presence of a metal halide e.g. copper chloride or mercuric

-16-

chlorides, and finally removing the Boc protecting groups by known methods, such as gaseous HCl in organic solvents.

5 This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully
10 in the claims which follow thereafter.

Experimental Details

Many of the specific compounds are identified by a
15 four-digit number from 2709 to 2732 and 3042, 3043 and 3069. Structures and names of these compounds may be found immediately following Chemical Synthesis Example 8.

20 CHEMICAL SYNTHESIS EXAMPLES

EXAMPLE 1

Preparation of phthalimidoalkylaminoindans [6].


25 A mixture of 0.14mol of aminoindan or IA, 0.14mol of bromoalkylphthalimide and 0.14 mol of K_2CO_3 in 180ml of MeCN was stirred at bath temperature 50°C for 8 hours. After cooling and filtration, the mother liquor was
30 evaporated. The residue was chromatographed on Keiselgel. Eluent: hexane-ethyl acetate 1:1.

In this way the following compounds were prepared;
35

-17-

Compounds of the
formula 1A

5



m=3, 56% yield.

10

$^1\text{H-NMR}(\text{CDCl}_3)$, delta, ppm: 7.85-7.63 (m, 4H), 7.31-7.10 (m, 4H), 4.2 (t, $J=6.5\text{Hz}$, 1H), 3.83-3.81 (m, 2H), 3.01-2.95 (m, 1H), 2.83-2.71 (m, 3H), 2.40-2.33 (m, 1H), 1.94-1.56 (m, 3H).

15

m=4, 54% yield.


$^1\text{H-NMR}(\text{CDCl}_3)$, delta, ppm: 7.88-7.65 (m, 4H), 7.32-7.15 (m, 4H), 4.22 (t, $J=6.5\text{Hz}$, 1H), 3.79-3.68 (m, 2H), 3.02-2.95 (m, 1H), 2.85-2.73 (m, 3H), 2.43-2.34 (m, 1H), 1.87-1.22 (m, 5H),

20

and

25

Compounds of
formula 1A'



30

m=m'=4, 8% yield.

$^1\text{H-NMR}(\text{CDCl}_3)$, delta, ppm: 7.88-7.68 (m, 8H), 7.32-7.11 (m, 4H), 4.46 (t, $J=7.5\text{Hz}$, 1H), 3.68-3.62 (m, 4H), 2.90-2.65 (m, 2H), 2.48-2.33 (m, 4H), 2.10-1.85 (m, 2H), 1.80-1.45 (m, 8H),

35

m=3, m'=4, 26% yield using a 100% excess of bromopropylphthalimide.

-18-

$^1\text{H-NMR}(\text{CDCl}_3)$, delta, ppm: 7.83-7.66 (m, 8H), 7.33-7.10 (m, 4H), 4.50 (t, $\gamma=7.5\text{Hz}$, 1H), 3.86-3.65 (m, 4H), 2.90-2.75 (m, 2H), 2.51-2.40 (m, 4H), 2.10-1.51 (m, 8H),


5 EXAMPLE 2

Hydrazinolysis of phthalimidoaminoindans [1]


A mixture 3.5mmol of phthalimidoaminoindan (as
10 prepared in Example 1) and 60% excess of 64% $\text{N}_2\text{H}_4 \cdot \text{H}_2\text{O}$
in 40 ml of abs. ethanol was refluxed with stirring
for 1.5 hrs. After cooling and filtration the mother
liquor was evaporated and the substance was extracted
by reflux with 50ml of ethyl acetate. The solvent was
15 evaporated. The residue was dissolved in 15ml of
ether and gas HCl in ether (15ml pH<1) was added with
stirring. The residue was filtered.

In this way the following compounds were prepared;

20



25



$m=3$ (2709) recrystallized from ethyl acetate/ethanol
(1.6:1), 82% yield, mpt 153-154°C.

30 $^1\text{H-NMR}(\text{D}_2\text{O})$, delta, ppm: 7.60-7.39 (m, 4H), 4.91-4.82
(m, 1H), 3.28-3.02 (m, 6H), 2.67-2.58 (m, 1H), 2.34-2.26
(m, 1H); 2.17-2.10 (m, 2H).

IR (KBr, disc), cm^{-1} 750, 1606, 1512, 3000-2500, 1444),
2700-2400, 3250.

35 m/z 190MH⁺ (free base)

Anal. Calc. for $\text{C}_{12}\text{H}_{20}\text{Cl}_2\text{N}_2$, C 54.75%, H 7.00%, Cl
26.94%, N 10.65%


-19-

Found. C 55.06%, H 7.82%, Cl 26.00%, N 11.31%

m=4 (2710) recrystallized from ethyl acetate/ethanol (1.6:1), 83% yield, mpt 214-216°C(decomp).

¹H-NMR(D₂O), delta, ppm: 7.56-7.34 (m,4H), 4.86-4.81 (m,1H), 3.21-2.98 (m,6H), 2.60-2.53 (m,1H), 2.30-2.23 (m,1H), 1.82-1.73 (m,4H).
m/z 205MH⁺ (free base).

and compounds;



m=m'=4 (2711)

was prepared by dissolution in water and stirring with Norit. After filtration and evaporation of water the residue was triturated with ether and crystallized from iso-propanol-ethanol 2:1, 45% yield.

¹H-NMR(D₂O), delta, ppm: 7.55-7.37 (m,4H), 5.24-5.21 (m,1H), 3.20-2.89 (m,10H), 2.57-2.51 (m,1H), 2.39-2.32 (m,1H), 1.90-1.56 (m,8H).

IR (KBr, disc), cm⁻¹ 770-735, 1480- 1440: 3000-2800, 1600,1500, 3000, 2500,2000, 1600-1575

m/z 275MH⁺ (free base)

Anal. Calc. for C₁₇H₃₂Cl₃N₃, C 53.20%, H 8.40%, Cl 27.45%, N 10.95%

Found. C 53.01%, H 8.08%, Cl 28.84%, N 11.16%.

m=m'=4 (R)-enantiomer of 2711, (2726)

The above procedure was repeated using (R)-aminoindan as the starting material and recrystallizing from ethanol-isopropanol. Yield 57%, mpt > 280°C.

-20-

$^1\text{H-NMR}(\text{D}_2\text{O})$, delta, ppm: 7.65-7.40 (m, 4H), 5.25 (m, 1H), 3.35-2.90 (m, 10H), 2.57 (m, 1H), 2.41 (m, 1H), 2.05-1.60 (m, 8H).

IR (KBr, disc), cm^{-1} 3000-2850, 2046, 2010, 1616, 1606, 1532, 1512, 1470, 1446, 1017, 974, 775, 747.

m/z 276 (MH^+ , 100, free base)

Anal. Calc. for $\text{C}_{17}\text{H}_{32}\text{Cl}_3\text{N}_3$, C 53.06%, H 8.38%, Cl 27.64%, N 10.92%

Found. C 52.35%, H 8.46%, N 10.85%.

10 $[\alpha]_D = -6.0^\circ(\text{H}_2\text{O})$

m=m'=4, (S)-enantiomer of 2711, (2727)

The above procedure was repeated using (S)-aminoindan as the starting material. Yield 56%, mpt $> 280^\circ\text{C}$.

15 Spectral properties were identical to 7262 above.
 $[\alpha]_D = +6.4^\circ(\text{H}_2\text{O})$

Similarly, the enantiomers of other compounds of general formula I may be made by using the corresponding enantiomeric 1-aminoindan as the starting compound.

25 m=3, m'=4 (2722) 45% yield
was prepared by dissolution in water and stirring with Norit. After filtration and evaporation of water the residue was triturated with ether and crystallized from iso-propanol-ethanol 2:1, 45% yield, mpt $227-228^\circ\text{C}$.

30 $^1\text{H-NMR}(\text{D}_2\text{O})$, delta, ppm: 7.56-7.39 (m, 4H), 5.28-5.26 (m, 1H), 3.32-2.97 (m, 10H), 2.60-2.54 (m, 1H), 2.40-1.62 (m, 7H).

IR (KBr, disc), V , cm^{-1} 770-735, 1480-1440: 3200-2800, 1600, 1500, 3000, 2500, 2000, 1600-1575.


35 m/z 261 MH^+ (free base)

Anal. Calc. for $\text{C}_{16}\text{H}_{30}\text{Cl}_3\text{N}_3$, C 51.82%, H 8.16%, Cl 28.69%, N 11.33%

-21-

Found. C 51.34%, H 7.77%, Cl 29.85%, N 11.63%.

and compounds



10 $m=m'=4$ (2728)

The above procedure was repeated using 1-aminotetralin as the starting material.

15 $^1\text{H-NMR}(\text{D}_2\text{O})$, delta, ppm: 7.53 (m, 1H), 7.48-7.33 (m, 3H), 5.04 (t, 1H), 3.50-2.65 (m, 10H), 2.35 (m, 1H), 2.15-1.40 (m, 11H).

IR (KBr, disc), V, cm^{-1} 3000-2850, 2014, 1616, 1600, 1534, 1437, 1408, 1012, 773, 754.

m/z 290 (MH^+ , 100, (free base)

20 Anal. Calc. for $\text{C}_{18}\text{H}_{24}\text{Cl}_3\text{N}_3$, C 54.21%, H 8.59%, Cl 26.67%, N 10.54%

Found. C 54.33%, H 8.84%, Cl 26.17%, N 10.65%.

25 EXAMPLE 3


3A. Protection of secondary amines with BOC_2O [1].

30 A solution of 16mmol of a secondary amine of Formula IA or IB and 24mmol of BOC_2O in 50ml of dichloromethane was stirred at R.T. for 3 hours. The reaction mixture was washed with 5% NaOH, dried, and evaporated. The residue was purified by column chromatography on Kieselgel. Eluent: hexane-ethyl acetate 2:1. Yield 85-95%.

35

In this manner the following compounds were prepared

Compounds of
Formula II



m=3 or m=4.

m=3;

1H -NMR ($CDCl_3$), delta, ppm: 7.82-7.70 (m, 4H), 7.14-7.05 (m, 4H), 5.73 and 5.25 (broad s 1H), 3.65-1.28 (m, 19H).

m=4:

1H -NMR ($CDCl_3$), delta, ppm: 7.81-7.70 (m, 4H), 7.13-7.00 (m, 4H), 5.72 and 5.25 (broad s 1H), 3.65-1.27 (m, 21H).


3B. Hydrazinolysis of BOC phthalimidoaminoindans [1].

A mixture of 50mmol of BOC-phthalimidoaminoindan of Formula IIA and 65mmol of 64% $N_2H_4 \cdot H_2O$ in 300ml of absolute ethanol was stirred at bath temperature 50°C for 10 hours. After cooling and filtration the mother liquor was evaporated. The residue was distributed in a mixture of ether-water. The ether phase was dried and evaporated, the residue was purified on Kieselgel. Eluent: dichloromethane-methanol-conc. ammonia 65:10:1. Yield 70-80%.

-23-


In this manner the following compounds were prepared;

5



Compounds of
Formula III

m=3;



¹H-NMR(CDCl₃), delta, ppm: 7.21-7.13 (m, 4H), 5.78 and
10 5.27 (broad s, 1H), 3.28-1.21 (m, 19H).

m=4:

¹H-NMR(CDCl₃), delta, ppm: 7.20-7.14 (m, 4H), 5.77 and
15 5.27 (broad s, 1H), 3.19-1.20 (m, 21H).


3C. Alkylation of primary amines.

A mixture of 41.4mmol of a primary amine of Formula
20 III, 82.8mmol of Cl(CH₂)_{m-1}CN and 124mmol of K₂CO₃ in
120ml of acetonitrile was refluxed with stirring for 3
hours, then an additional 41.4mmol of Cl(CH₂)_{m-1}CN was
added, and the mixture was refluxed for 4 hours.
After addition of a further 82.8mmol of Cl(CH₂)_{m-1}CN the
25 reaction mixture was refluxed for 3 hours. After
cooling, filtration and evaporation, the residue was
purified by column chromatography on Kieselgel.
Eluent: dichloromethane-methanol 95:5. Yield 60-65%. m
is 3 or 4 depending on the compound being synthesized.


30

In this manner the following compounds were prepared;

35



Compounds of
Formula IV



-24-

m=3, m'=4

¹H-NMR(CDCl₃), delta, ppm: 7.21-7.15 (m, 4H), 5.70 and 5.30 (broad s, 1H), 2.98-1.25 (m, 25H).


5 m/z 258MH⁺

3D. Protection of secondary amines with BOC₂O.


10 A solution of 16mmol of a secondary amine of Formula IVA or IVB and 24mmol of BOC₂O in 50ml of dichloromethane was stirred at R.T. for 3 hours. The reaction mixture was washed with 5% NaOH, dried, and evaporated. The residue was purified by column chromatography on Kieselgel. Eluent: hexane-ethyl acetate 2:1. Yield 85-95%.

In this manner the following compounds were prepared;

20



Compounds of
Formula V



m=3, m'=4

25 ¹H-NMR(CDCl₃), delta, ppm: 7.21-7.16 (m, 4H), 5.78 and 5.28 (broad s, 1H), 3.10-1.24 (m, 34H).

3E. Reduction of nitriles with LiAlH₄

30

To a suspension 4.0g of LiAlH₄ in 150ml of ether 5.8mmol a nitrile of Formula VB in 50ml of ether was added with stirring in an ice-bath, the suspension was then stirred without ice-bath for 0.5hrs. Excess of
35 LiAlH₄ was destroyed followed by addition of 4.0ml of water, 4.0ml of 10% NaOH and 12.0ml of water with cooling. The organic phase was decanted and additional


-25-

amount of the substance was extracted by dichloromethane. The combined organic phases were washed, dried and evaporated. The residue was purified by column chromatography on Kieselgel.


5 Eluent: dichloromethane-methanol - conc. ammonia 65:10:1. Yield 60-70%.

In this manner the following compounds were prepared;

10



Compounds of
Formula VI



15 $m=3, m'=4$



$^1\text{H-NMR}(\text{CDCl}_3)$, delta, ppm: 7.22-7.14 (m, 4H), 5.78 and 5.30 (broad s, 1H), 3.04-1.26 (m, 36H).

20 3F. Deprotection of N-BOC derivatives.

An excess of HCl in ether (18ml, pH<1) was added to a solution 2.5mmol of an N-BOC derivative of Formula VI in 6ml of methanol. The mixture was stirred at R.T.

25 for 4 hours. The precipitate was collected by filtration and washed with ether.

30 In this manner the following compounds were prepared;



35

-26-

m=3,m'=4 yield 50% mpt 239-240°C (2720).

¹H-NMR(D₂O), delta, ppm: 7.56, 7.34(m, 4H), 4.87-4.83 (m,1H), 3.23-2.98 (m,10H), 2.63-2.53 (m,1H), 2.30-2.22 (m,1H), 2.15-2.08 (m, 2H), 1.82-1.71 (m, 4H).

5 IR (KBr, disc), cm⁻¹ 770-735, 1600, 1500, 2000, 2300-2700, 2500, 3000, 3400, 3500.

m/z 262 (MH⁺, 100, free base)

Anal. Calc. for C₁₆H₃₀Cl₃N₃ 0.7H₂O*, C 50.12%, H 8.18%, Cl 27.74%, N 10.96%

10 Found. C 50.16%, H 7.98%, Cl 28.09%, N 11.17%.

m=4,m'=3 (2724) 45% yield, mpt 237-238°C

¹H-NMR(D₂O), delta, ppm: 7.67-7.40 (m,4H), 4.90-4.86 (m,1H), 3.21-3.01 (m,10H), 2.71-2.56 (m,1H), 2.33-2.25 (m,1H), 2.17-2.09 (m,2H), 1.82-1.34 (m,4H).

15 IR (KBr, disc), cm⁻¹ 750, 1608,1500, 2700-2250, 2943, 2500, 2042

m/z 262MH⁺ (free base)

20 Anal. Calc. for C₁₆H₃₀Cl₃N₃ 0.7H₂O* C 50.12%, H 8.18%, Cl 27.74%, N 10.96%,

Found. C 49.10%, H 7.81%, Cl 27.68%, N 11.20%.

* water content determined by Karl Fischer analysis.

25

EXAMPLE 4

4A. Alkylation of primary amines.

30 A mixture of 41.4mmol of a primary amine of Formula VI, 82.8mmol of Cl(CH₂)_{m-1}CN and 124mmol of K₂CO₃ in 120ml of acetonitrile was refluxed with stirring for 3 hours, then an additional 41.4mmol of Cl(CH₂)_{m-1}CN was added, and the mixture was refluxed for 4 hours.


35 After addition of a further 82.8mmol of Cl(CH₂)_nCN the reaction mixture was refluxed for 3 hours. After cooling, filtration and evaporation the residue was

-27-

purified by column chromatography on Kieselgel.

Eluent: dichloromethane-methanol 95:5. Yield 60-65%. m is 3 or 4 depending on the compound being synthesized.

5 In this manner the following compounds were prepared;




4B. Protection of secondary amines with BOC₂O [1].

15 A solution of 16mmol of a secondary amine of Formula VIIA or VIIB and 24mmol of BOC₂O in 50ml of dichloromethane was stirred at R.T. for 3 hours. The reaction mixture was washed with 5% NaOH, dried, and evaporated. The residue was purified by column

20 chromatography on Kieselgel. Eluent: hexane-ethyl acetate 2:1. Yield 85-95%.

In this manner the following compounds were prepared;



4C. Reduction of nitriles with LiAlH₄

35 To a suspension 4.0g of LiAlH₄ in 150ml of ether 5.8mmol a nitrile of Formula VIIIB in 50ml of ether was added with stirring in an ice-bath, the suspension was then stirred without ice-bath for 0.5hrs. Excess


-28-

of LiAlH_4 was destroyed followed by addition of 4.0ml of water, 4.0ml of 10% NaOH and 12.0ml of water with cooling. The organic phase was decanted and additional amount of the substance was extracted by
 5 dichloromethane. The combined organic phases were washed, dried and evaporated. The residue was purified by column chromatography on Kieselgel. Eluent: dichloromethane-methanol - conc. ammonia 65:10:1. Yield 60-70%


10

In this manner the following compounds were prepared;

15



Compounds of
Formula VIII



4D. Deprotection of N-BOC derivatives.



20

An excess of HCl in ether (18ml, $\text{pH} < 1$) was added to a solution 2.5mmol of an N-BOC derivative of Formula VIII in 6ml of ether. The mixture was stirred at R.T. for 4 hours. The precipitate was collected by
 25 filtration and washed with ether. The salt was stirred in water with Norit. After filtration and evaporation the residue was triturated with methanol, filtered and washed with ether.

30

In this manner the following compounds were prepared;

5



10

$m=3, m'=4$ (2721) mpt $>280^\circ\text{C}$.

$^1\text{H-NMR}(\text{D}_2\text{O})$, delta, ppm: 7.56-7.35 (m, 4H), 4.88-4.85 (m, 1H), 3.24-2.98 (m, 14H), 2.63-2.54 (m, 1H), 2.30-2.33 (m, 1H), 2.16-2.06 (m, 4H), 1.79-1.77 (m, 4H).

m/z 319319MH $^+$

15

$m=4, m'=3$ (2725) mpt $>280^\circ\text{C}$.

$^1\text{H-NMR}(\text{D}_2\text{O})$, delta, ppm 7.56-7.34 (m, 4H), 4.86-4.83 (m, 1H), 3.21-3.02 (m, 14H), 2.61-2.53 (m, 1H), 2.30-2.22 (m, 1H), 2.16-2.19 (m, 2H), 1.81-1.73 (m, 8H).

20

IR (KBr, disc), cm^{-1} 750, 1610, 1500, 2700-2250 (N $^{\circ}$ HR), 3000, 2500, 2000 (NH $_3$ $^+$)

m/z 333MH $^+$ (free base)

Anal. Calc. for $\text{C}_{20}\text{H}_{40}\text{Cl}_4\text{N}_4 \cdot 0.5\text{H}_2\text{O}$ C 49.20%, H 8.20%, Cl 29.10%, N 11.50%,

25

Found. C 47.82%, H 8.39%, Cl 29.52%, N 11.99%.

* water content determined by Karl Fischer analysis.

30

EXAMPLE 5

A. 1,3-Indandione Dioxime

35

A mixture of 75 g of potassium carbonate, 30 g of hydroxylamine hydrochloride, 15 g of 1,3-indandione and 300 mL of 95% ethanol was heated to reflux with stirring for 15 min at which time it was poured into

-30-

400 mL of ice. The resulting precipitate was collected by filtration, washed with cold water and dried in vacuo to give 17.9 g of the title compound as an amorphous solid. NMR (CDCl₃, CD₃OD): 7.42-7.76 (m, 4H); 3.36 (s, 2H).

B. 1,3 Diaminoindan (cis/trans)

To a solution of 20 g of a 1,3-indandione dioxime in 2 L of absolute ethanol was cautiously added 160 g of sodium in portions during 1 hr (at such a rate to maintain reflux). After disappearance of the sodium, 300 mL of water was slowly added and the solution concentrated in vacuo to remove most of the ethanol. The resulting solution was acidified with conc. hydrochloric acid to pH 1-2 followed by extraction with methylene chloride to recover unreacted starting material. The aqueous layer was basified with 10% sodium hydroxide to pH 12 and extracted with methylene chloride which gave 11.7 grams of crude 1,3-diaminoindan upon solvent removal in vacuo. Purification of this material by flash column chromatography eluting with 95% ethanol/ammonium hydroxide (400:16) gave 8.75 g of the title compound as an approximately 60/40 mixture of cis to trans isomers.

C. cis-N,N'-Diacetyl-1,3-diaminoindan (2717)

To a solution of 17.55 g of cis/trans-1,3-diaminoindan in 102 mL of pyridine was slowly added 102 mL of acetic anhydride. After 17 hrs the reaction mixture was added to 250 mL of water and 300 mL of methylene chloride. The insoluble precipitate was isolated from this three-phase mixture by filtration and dried to

-31-

give 8.7 g of crude title compound which was recrystallized from ethanol to give 5.4 g pure material.

5 ¹NMR (CDCl₃, CD₃OD) delta: 7.3 (s, 4H); 5.3 (m, 2H); 2.0 (s, 6H); 2.9 and 1.7 (m, 2H).

D. trans-N,N'-Diacetyl-1,3-diaminoindan (2718)

10 The combined methylene chloride extracts from the above example were concentrated in vacuo, suspended in 100 mL of water and basified with 10% sodium hydroxide to pH 12. Evaporation of the repeated methylene chloride extracts gave 9.4 g of crude N,N'-diacetyl-1,3-diaminoindan as a 35:65 cis/trans mixture. The
15 title compound (3.5 g) was isolated by flash column chromatography eluting with ethyl acetate/ethanol (95:5).

¹NMR (CDCl₃, CD₃OD) delta: 7.35 (m, 4H); 5.59 (m, 2H); 2.33 (t, 2H); 2.02 (s, 6H).

20

E. cis-1,3-Diaminoindan hydrochloride (2715)

A solution of 100 mg of cis-N,N'-diacetyl-1,3-diaminoindan in 4 mL of 5% hydrochloric acid was
25 heated at reflux for 6 hrs at which time it was cooled and 10% sodium hydroxide was added to pH 12-13. The combined methylene chloride extracts were evaporated to dryness to give 65 mg of crude diamine which was treated with methylene chloride and etheric hydrogen
30 chloride to give the title compound (35 mg).

¹NMR (CD₃OD) delta: 7.48 (m, 4H); 4.58 (d, d, 2H); 1.74, 3.05 (m, 2H).

F. trans-1,3-Diaminoindan hydrochloride (2716)

35

The title compound was prepared according to the method of the above example 5E, except that trans-

-32-

N,N'-diacetyl-1,3-aminoindan was used as starting material.

¹NMR (CD₃OD) delta: 7.35 (m, 4H); 4.55 (t, 2H), 2.15 (t, 2H).

5

EXAMPLE 6

6A. Alkylation of 1,3-diaminoindan dihydrochloride


10

A mixture 14.9mmol of 1,3-diaminoindan dihydrochloride, 42.0mmol of 4-chlorobutyronitrile and 84.0mmol of K₂CO₃ in 50ml of acetonitrile was refluxed for 3 hrs. Then an additional 84.0mmol of 4-chlorobutyronitrile was added, and the mixture was refluxed for 4 hrs. After cooling and filtration the mother liquor was evaporated. The residue and 42.0mmol of BOC₂O were dissolved in 150ml of dichloromethane and stirred at R.T. overnight. The reaction mixture was washed with 5% NaOH, dried, and evaporated. The residue was purified by column chromatography on Kieselgel, eluting with hexane-ethyl acetate (2:1) to isolate the BOC-mono-alkylated (XIII) and di-alkylated (XIV) products.

20

In this manner the following compounds were prepared;

30



Compounds of Formulae
XIII and XIV

cis R=BOC, R'=CH₂CH₂CH₂CN, R''=H: IR (neat): 2245cm⁻¹

35

-33-

cis R=BOC, R'=R''=CH₂CH₂CH₂CN: IR (neat): 2245cm⁻¹

5 trans R=BOC, R'=R''=CH₂CH₂CH₂CN: IR (neat): 2245cm⁻¹


10 6B. Reduction of nitriles with LiAlH₄

To a suspension 4.0g of LiAlH₄ in 150ml of ether
5.8mmol a nitrile of Formula XIII or XIV in 50ml of
ether was added with stirring in an ice-bath, the
suspension was then stirred without ice-bath for
15 0.5hrs. Excess LiAlH₄ was destroyed followed by
addition of 4.0ml of water, 4.0ml of 10% NaOH and
12.0ml of water with cooling. The organic phase was
decanted and an additional amount of the substance was
extracted by dichloromethane. The combined organic
20 phase was washed, dried and evaporated. The residue
was purified by column chromatography on Kieselgel.
Eluent: dichloromethane-methanol - conc. ammonia
65:10:1.

25

In this manner the following compounds were prepared;

30



cis R=BOC, R'=CH₂CH₂CH₂CH₂NH₂, R''=H:

35

-34-

cis R=BOC, R'=R''=CH₂CH₂CH₂CH₂NH₂:

m/z 491MH⁺

5

trans R=BOC, R'=R''=CH₂CH₂CH₂CH₂NH₂:

6C. cis-N-(4-aminobutyl)-1,3-diaminoindan
trihydrochloride (2731):

10

A solution of the above di-BOC derivative in methanol/ether was treated with gaseous hydrogen chloride for 6 hrs at RT. The precipitate was isolated by filtration and washed with ether.

15

A solution of this crude title compound (0.22g) in 15ml of methylene chloride/water (2:1) was again treated with BOC anhydride (0.62g) and 0.28g of potassium carbonate at 50°C for 20 min. The organic phase was separated, washed with 5% sodium hydroxide and evaporated to dryness. Flash column chromatography on silica eluting with hexane/ethyl acetate (2:1) gave 170mg of the tri-BOC derivative. Acidolysis was accomplished with 1ml of trifluoroacetic acid at RT during 20 min. After evaporation of the trifluoroacetic acid, the title compound (41mg) was isolated from methanol/ether after treatment with hydrogen chloride gas.

20

25

30

¹H-NMR(CDCl₃), delta, ppm: 7.85-7.80 (m,1H), 7.68-7.56 (m,3H), 4.97-4.89 (m,2H), 3.30-3.18 (m,3H), 3.03-2.98 (m,2H), 2.43-2.37 (m,1H), 1.98-1.77 (m,4H).
m/z 220MH⁺ (free base)

6D. trans-N,N'-(Bis-4-aminobutyl)-1,3-diaminoindan
tetrahydrochloride (2732):

35

A solution of the above di-BOC derivative in methylene

-35-

chloride was treated with BOC-anhydride at RT for 3hrs at which time the reaction mixture was washed with 5% sodium hydroxide and concentrated to dryness. The residue was purified by flash column chromatography on silica eluting with hexane/ethyl acetate (2:1) to give the tetra-BOC derivative. Acidolysis was accomplished with 1ml of trifluoroacetic acid at RT during 20 min. After evaporation of the trifluoroacetic acid, the title compound was isolated from methanol/ether after treatment with hydrogen chloride gas.

¹H-NMR(CDCl₃), δ , ppm: 7.90-7.87 (m, 2H), 7.62-7.59 (m, 2H), 5.23-5.20 (m, 2H), 3.30-3.15 (m, 4H), 3.02-2.98 (m, 4H), 2.91-2.88 (m, 2H), 1.96-1.76 (m, 8H).
m/z 291MH⁺

EXAMPLE 7:

Preparation of (XVIII)

A mixture of racemic aminoindan (12.38g, 0.093 mol), potassium carbonate (12.84g, 0.093 mol), N-(3-Bromopropyl)-phthalimide (24.92g, 0.093 mol) and acetonitrile (120mL) was stirred and heated at 50-55°C under nitrogen for 10 hours. The mixture was then cooled and filtered under vacuum. The mother liquor was evaporated at reduced pressure to give 30.4g of crude product which was purified by flash chromatography to give 21.27g (71.5% yield) of an off-white solid.

Preparation of (XIX)

A mixture of XVIII (17.45g, 0.055 mol), 4-bromobutyronitrile (8.1g, 0.055 mol), potassium carbonate (30.19g, 0.218 mol) and acetonitrile (440mL) was stirred and heated at reflux under nitrogen for 5 hours. The mixture was cooled to 60°C and additional 4-bromobutyronitrile (32.43g, 0.219 mol) was added. The mixture was now stirred and heated at reflux under

-36-

nitrogen for 20 hours. The mixture was then cooled and filtered and the mother liquor was evaporated under vacuum. The residue was heated under high vacuum to remove the excess 4-bromobutyronitrile and the residual viscous oil was purified by flash chromatography to give 17.80g (84% yield).

¹H NMR δ (DMSO): 7.83 (s, 4H, Ph), 7.28-7.05 (m, 4H, Ph), 4.46 (t, 1H, Cl-H), 3.67 (m, 1H, C3-H), 3.47 (m, 1H, C3-H'), 2.75 (m, 2H), 2.60-2.30 (m, 6H), 2.05 (m, 1H), 1.95 - 1.60 (m, 5H) ppm.

Preparation of (XX)

A mixture of XIX (17.45g, 0.045 mol), 55% hydrazine hydrate (4.25g, 0.073 mol, 60% excess) and absolute ethanol (550mL) was stirred and heated at reflux under nitrogen for 4 hours. On cooling a white solid precipitated; it was filtered and washed with cold ethanol (3 x 75mL). The combined filtrate was evaporated under vacuum to give a crude yellow solid. This crude solid was dissolved in a mixture of saturated brine (200mL) and hot ethyl acetate (300mL) and the layers were separated. The aqueous layer was reextracted with ethyl acetate (3 x 100mL). The combined organic layer was dried (Na₂SO₄) and solvent was removed at reduced pressure to give 10.8g of a yellow oil. The crude product was purified by flash chromatography to give 7.65g (66% yield) of a yellow oil.

¹H NMR δ (DMSO): 7.30-7.10 (m, 4H, Ph), 4.45 (t, 1H, Cl-H), 2.95-2.65 (m, 2H), 2.60-2.30 (m, 8H), 2.05 (m, 1H), 1.86 (m, 1H), 1.70 (m, 2H), 1.50 (m, 2H) ppm
MS: 259 (NH⁺, 100)

Preparation of (XXI)

A solution of XX (7.65g, 0.030 mol) and di-tert-butyl dicarbonate (9.80g, 0.045 mol) in methylene chloride

-37-

(400mL) was stirred at 20°C under nitrogen for 8 hours. The reaction mixture was then washed with 5% aqueous NaOH solution (150mL) and the layers were separated. The aqueous layer was reextracted with methylene chloride (3 x 75mL) and the combined layer was dried and evaporated to 14.5 grams of a yellow oil. The crude product was purified by flash chromatography to give 10.15g (95% yield) of a yellow oil.

Preparation of (XXII)

To a suspension of LAH (19.35g, 0.51 mol) in anhydrous ether (730mL) was added IV (10.0g, 0.028 mol) dissolved in anhydrous ether (230mL). The addition was carried out at 0-5°C with stirring under a nitrogen atmosphere. After the addition was complete, the suspension was stirred at 5°C for an additional 30 minutes and then it was stirred at 25°C under nitrogen for 3 hours. The mixture was now cooled to 5°C and the excess LAH was decomposed by the addition of water (110mL), 10% sodium hydroxide solution (320mL) and more water (75mL). Ethyl acetate (365mL) was then added to the stirred mixture and it was filtered through a celite pad. The pad was washed with ethyl acetate (3 x 150mL) and the layers were separated. The aqueous layer was reextracted with ethyl acetate (150mL). The combined ethyl acetate layer was dried (Na_2SO_4) and solvent was removed under vacuum to give 9.27g (92% yield) of a viscous yellow oil.

Preparation of (XXIII)

A mixture of XXII (7.9g, 0.022 mol), potassium carbonate (3.03g, 0.022 mol), N-(3-Bromopropyl)-phthalimide (5.89g, 0.022 mol) and acetonitrile (175mL) was stirred and heated at 55-60°C under nitrogen for 7 hours. The mixture was then cooled and filtered under vacuum. The mother liquor

-38-

was evaporated at reduced pressure to give 12.3g of a viscous oil which was purified by flash chromatography to give 3.3g (28% yield) of a white solid.

5 Preparation of (XXIV)

A mixture of XXIII (3.0g, 5.47 mmol), 55% hydrazine hydrate (0.51g, 8.77 mmol, 60% excess) and absolute ethanol (120mL) was stirred and heated at reflux under nitrogen for 5 hours. It was then evaporated
10 under vacuum to give a white solid which was dissolved in a mixture of saturated brine (70mL) and hot ethyl acetate (100mL). The layers were separated and the aqueous layer was reextracted with ethyl acetate (2 x 70mL). The combined organic layer was dried (Na_2SO_4)
15 and solvent was removed at reduced pressure to yield 2.25g (98% yield) of a viscous yellow semi-solid.

Preparation of (XXV)

A solution of XXIV (2.25g, 5.38 mmol) and
20 di-tert-butyl dicarbonate (4.46g, 20.4 mmol) in methylene chloride (75mL) was stirred at 25°C under nitrogen for 22 hours. The reaction mixture was then washed with 5% aqueous NaOH solution (100mL) and the layers were separated. The aqueous layer was
25 reextracted with methylene chloride (3 x 40mL) and the combined organic layer was dried (Na_2SO_4). Removal of solvent at reduced pressure gave 5.65 grams of a yellow oil which was purified by flash column chromatography to give 3.26g (98% yield) of a yellow
30 oil.

Preparation of 3069

Ether saturated with HCl gas (30mL) was added to a solution of XXV (3.16g, 5.22 mmol) dissolved in
35 methanol (60mL). The solution was stirred at 25°C for 3 1/2 hours. Solvents were removed under vacuum to give 2.29g of an off-white solid which was purified by

-39-

heating it in water (300mL) containing Norit (480mg) at 90-95°C for 5 minutes. The mixture was filtered hot through a celite pad and the pad was washed with hot water (3 x 150mL). The colorless filtrate was
5 evaporated under vacuum at 60°C. Hexane was added to the residue and solvent was removed under vacuum. Finally, methanol (100mL) was added to the residue. Its removal under reduced pressure gave 2.16g (91%) of a hygroscopic white solid.

10

Anal. calcd for $C_{19}H_{38}Cl_4N_4 \cdot H_2O$: C, 47.31; H, 8.36; Cl, 29.40; N, 11.62
Found: C, 47.71; H, 8.21; Cl, 29.25; N, 11.65

15

1H NMR δ (DMSO): 7.62 (d, 1H, Ph), 7.58-7.40 (m, 3H, Ph), 5.28 (dd, 1H, Cl-H), 3.30-3.0 (m, 14H,), 2.58 (m, 1H, C2-H'), 2.40 (m, 1H, C2-H), 2.30-2.05 (m, 4H), 2.0-1.6 (m, 4H) ppm

20

MS : 319 (MH⁺)

HPLC: 99%

EXAMPLE 8:

25

Preparation of II

A mixture of N,N-Bis-4-aminobutyl racemic aminoindan (compound no. 2711, 2.5g, 6.5mmol), N,N-di-tert-butoxycarbonylthiourea, (Iwanowicz E et al. Synthetic
30 Communications (1993) 23, 1443), (3.5g, 13.0 mmol), triethyl amine (3.61g, 35.75 mmol) and dry DMF (19.5mL) was stirred at 0°C under nitrogen for an hour. The ice bath was removed and the mixture was
35 then stirred at 22-24°C under nitrogen for 22 hours. Ethyl acetate (200mL) was added and the mixture was filtered through a Celite pad. The pad was washed

-40-

with ethyl acetate (4 x 70mL). The combined filtrates were washed with water (200mL), sat. sodium chloride (200mL) and dried (Na_2SO_4). Removal of solvent gave 6.2g of a viscous semi-solid. The crude product was purified by flash chromatography using hexane/ethyl acetate 70/30 as eluent. This gave 3.5g (71% yield) of an off-white hygroscopic solid.

Preparation of III

A mixture of N-4-aminobutyl racemic aminoindan (2710, 3.33g, 12.0 mmol), N,N-di-tert-butoxycarbonylthiourea, (Iwanowicz E et al. Synthetic Communications (1993) 23, 1443), (3.32g, 12.0 mmol), triethylamine (4.85g, 48.0 mmol) and dry DMF (30mL) was treated and worked up according to the conditions used for the preparation of II. Removal of solvent gave 6.0g of a viscous oil/semi-solid. The crude product was purified by flash chromatography using methylene chloride/methanol 94/6 as eluent. This gave 3.5g (65.5% yield) of a viscous brown oil.

Preparation of (N,N-di-(4-Guanidinobutyl)-aminoindan·3HCl (3042)

To a solution of II (3.0g, 3.95 mmol) dissolved in ether (10mL) was added ether saturated with HCl gas (10mL). A gummy oil precipitated. Then methanol (40mL) was added and the solution was stirred at 25°C for 30 minutes. Solvents were removed under vacuum to give 3.2g of a white solid. It was purified by heating it in water (150mL) containing Norit (500mg) at 90-95°C for 5 minutes. The mixture was filtered hot through a Celite pad and the pad was washed with hot water (2 x 100mL). The colorless filtrate was evaporated under vacuum at 60°C. Hexane was added to the residue and solvent was removed under vacuum.

-41-

Finally, methanol (100mL) was added to the residue. Its removal under reduced pressure gave a white solid (1.60g, 86%) which was dried at 40°C under high vacuum and stored in a desiccator over phosphorus pentoxide.

5

Anal. for $C_{19}H_{36}C_{13}N_7 \cdot 2H_2O$: C, 45.2; H, 7.98, N, 19.42

Found: C, 44.90, H, 6.65; N, 19.06

1H NMR δ (D_2O): 7.60-7.30 (m, 4H, Ph), 5.20 (dd, 1H, C1-H)

3.40-3.0 (m, 9H), 2.90 (m, 1H, C3-H), 2.52 (m, 1H, C2-

10 H), 2.36 (m, 1H, C2-H¹), 2.0-1.4 (m, 8H) ppm.

FABMS: 360 (MH⁺), 396 (MH⁺+HCl).

IR (KBr): 3397, 1665, 1464, 1167, 766 cm^{-1} .

15

HPLC: 96%

Preparation of (N-(4-Guanidinobutyl)-aminoindan-2HCl)
(3043)

20

Ether saturated with HCl gas (15mL) was added to a solution of III (305g, 7.85 mmol) dissolved in ether (10mL). A gummy solid precipitated. Then methanol (50mL) was added and the solution was stirred at 25°C
25 for 30 minutes. Solvents were removed under vacuum to give 3.3g of a white solid, which was purified by heating in water (180mL) containing Norit (560mg) at 90-95°C for 5 minutes. The mixture was filtered hot through a Celite pad and the pad was washed with hot
30 water (2 x 100mL). The colorless filtrate was evaporated under vacuum at 60°C. Hexane was added to the residue and solvent was removed under vacuum. Finally, methanol (100mL) was added to the residue. Its removal under reduced pressure gave 2.14g (86%) of
35 an extremely hygroscopic white solid which was dried at 25°C under high vacuum and stored in a desiccator over P_2O_5 .

-42-

Anal. calcd. for $C_{14}H_{24}Cl_2N_4 \cdot H_2O$: C, 49.85; H, 7.77; N, 16.6

Found: C, 49.11; H, 7.63; N, 16.70








1H NMR δ (D_2O): 7.60-7.30 (m, 4H, Ph), 4.8 (m, 1H, Cl-H), 3.30-
5 3.05 (m, 5H), 3.0 (m, 1H, C3-H), 2.55 (m, 1H, C2-
H), 2.26 (m, 1H, C2-H) 1.80-1.60 (m, 4H) ppm







FABMS: 247 (MH^+), 283 ($MH^+ + HCl$).

10 IR (KBr): 3349, 3161, 2950, 2803, 1665, 1462, 1169, 765 cm^{-1} .


HPLC: 95%

STRUCTURES OF COMPOUNDS


- 2709 
 $\text{HN}-(\text{CH}_2)_3-\text{NH}_2 \cdot (\text{HCl})_2$
- 2710 
 $\text{HN}-(\text{CH}_2)_4-\text{NH}_2 \cdot (\text{HCl})_2$
- 2711 
 $\text{H}_2\text{N}-(\text{CH}_2)_4-\text{N}-(\text{CH}_2)_4-\text{NH}_2 \cdot (\text{HCl})_3$
- 2726 (R)-2711
- 2727 (S)-2711
- 2722 
 $\text{H}_2\text{N}-(\text{CH}_2)_4-\text{N}-(\text{CH}_2)_3-\text{NH}_2 \cdot (\text{HCl})_3$
- 2728 
 $\text{H}_2\text{N}-(\text{CH}_2)_4-\text{N}-(\text{CH}_2)_4-\text{NH}_2 \cdot (\text{HCl})_3$
- 2720 
 $\text{HN}-(\text{CH}_2)_3-\text{NH}-(\text{CH}_2)_4-\text{NH}_2 \cdot (\text{HCl})_3$
- 2724 
 $\text{HN}-(\text{CH}_2)_4-\text{NH}-(\text{CH}_2)_3-\text{NH}_2 \cdot (\text{HCl})_3$




3069



3042



3043



-46-

Names of Some Compounds

	<u>Compound Number</u>	<u>Compound Name</u>
5	2709	N-(3-aminopropyl)-1-aminoindan
	2710	N-(4-aminopropyl)-1-aminoindan
	2711	racemic N,N-di(4-amino-n-butyl)-1-aminoindan
10	2726	(R)-N,N-di(4-amino-n-butyl)-1-aminoindan
	2727	(S)-N,N-di(4-amino-n-butyl)-1-aminoindan
15	2722	N-(4-amino-n-butyl)-N-(3-amino-n-propyl)-1-aminoindan
	2728	N,N-di(4-amino-n-butyl)-1-amino-1,2,3,4-tetrahydronaphthalene
20	2720	N-[N-(4-aminobutyl)-3-aminopropyl]-1-aminoindan or N-(1-indanyl)-N'-(4-aminobutyl)-1,3-diaminopropane
25	2724	N-[N-(3-aminopropyl)-4-aminobutyl]-1-aminoindan or N-(1-indanyl)-N'-(3-aminopropyl)-1,4-diaminobutane
30	2721	N-{[N-(3-aminopropyl)-4-aminobutyl]-N'-3-aminopropyl}-1-aminoindan or N-(1-indanyl)-N'-[N-(3-aminopropyl)-4-aminobutyl]-1,3-diaminopropane
35	2725	N-{[N-(4-aminobutyl)-3-aminopropyl]-N'-4-aminobutyl}-1-aminoindan or N-(1-indanyl)-N'-[N-(4-aminobutyl)-3-aminopropyl]-1,4-diaminobutane

-47-

	2717	cis N,N'-diacetyl-1,3-diaminoindan
	2718	trans N,N'-diacetyl-1,3-diaminoindan
5	2715	cis 1,3-diaminoindan
	2716	trans 1,3-diaminoindan
	2731	N-(4-aminobutyl)-1,3-diaminoindan
10	2732	N,N'-di(4-aminobutyl)-1,3-diaminoindan
	3069	N-(3-aminopropyl)-N-[N-(3-aminopropyl)-4-aminobutyl]-1-aminoindan
15	3042	N,N-di(4-guanidinobutyl)-1-aminoindan
	3043	N-(4-guanidinobutyl)-1-aminoindan

BIOLOGICAL EXAMPLES

1. Cultures of dissociated neonatal rat cerebellum

5 A. Reversal of NMDA induced cell death.

The cerebellum was aseptically dissociated from 6 or 7-day old rat pups and placed in a 15ml sterile plastic conical tube containing 3ml of Dulbecco's modified Eagle's medium (DMEM) with a high glucose concentration (1g/ml) and 2mM (v/v) L-glutamine and an antibiotic antimetabolic mixture. The cerebella were then dissociated after 20-25 passages through a sterile 13 gauge, 10cm long stainless steel needle attached to a 5ml syringe with an inserted 45 micrometer pore nylon sieve. The dissociated cells were centrifuged at 200g for 5 minutes. The supernatant was discarded and the cells resuspended in medium enriched with 15% (v/v) heat inactivated fetal calf serum. Cell viability was determined by the trypan blue exclusion test.

Cells were plated at a density of 200/mm² on poly-L-lysine coated surfaces. Poly-L-lysine coated glass coverslips were prepared at least one hour in advance of plating by immersing sterile coverslips in sterile distilled water solution containing 15 microgram/ml poly-L-lysine, and washing in sterile water just prior to use. The plated cells were covered with enriched medium and incubated at 37°C in an atmosphere of 5% CO₂ in air and 97% humidity. After three days in culture, the media was replaced with media containing the desired test compound. Each test compound was tested in duplicate. Toxic-dose response was determined for each compound over a range from 0.005-1000 micromole.

Four groups were run in each set of experiments;

I. Control, consisting of enriched media alone,

II. N-methyl-D-aspartate (NMDA, 1mM for 3 hours) as

-49-

the cytotoxic challenge,

III. Test compound plus NMDA, and

IV. Positive control, spermine, spermidine or putrescine

5 (0.01 micromoles) plus NMDA.

Nerve cell survival was evaluated by phase contrast microscopy and tryptan blue staining after 24 hours.

Results

10 The results are shown in Table 1 below. Surviving cells in Culture are measured relative to control (100%) as described above. Percent protection is the Cell Survival for the test compound minus the NMDA effect. Thus, maximal protection is 100% minus 30%
15 NMDA effect i.e. 70%. The Effective Protection was calculated as the percent of the Percent Protection (X) divided by the maximal protection value (e.g. $X \times 100/70$).

Table 1

COMPOUND	EXPERMNT'L GROUP Dose (μ M)	SURVIVING CELL IN CULTURE	PERCENT PROTECTION	EFFECTIVE PROTECTION
	Control	100		
	NMDA	30		
	Max protection		70	100
	Spermine + NMDA	68	38	54
	Spermidine + NMDA	78	48	69
	Putrescine + NMDA	30	0	0
(R)-1- aminoindan	0.005	80	50	71
	0.010	30	0	0
	0.100	28	-2	-
2709	0.005	26	-4	-
	0.010	53	23	33
	0.100	59	29	41
	1.000	43	13	18*
	10.00	38	8	11
2710	0.100	72	42	60
	1.000	70	40	57
2711	0.005	74	44	63
	0.010	68	38	54
	0.100	62	32	46
2715	0.005	49	19	27
	0.100	42	12	17
	1.000	32	2	3
2716	0.005	65	35	50
	0.010	30	0	0
	0.100	25	-5	-
	1.000	30	0	0
2720	0.005	72	42	60
	0.010	38	8	11
	0.100	32	2	3
2721	0.005	28	-2	-
	0.010	26	-4	-
2722	0.005	115	85	121
	0.010	87	57	81
2724	0.005	95	65	93
	0.010	87	57	81

-51-

2725	0.005	72	42	60
	0.010	38	8	11
	0.100	32	2	3
2728	0.005	20	-10	-
	0.010	86	56	80
2726	0.005	129	99	141
	0.010	55	25	36
2727	0.005	30	0	0
	0.010	55	25	36

B: In Vitro Cell Toxicity

Toxic dose-response curves were compiled using the
5 protocol described above. The TD50 values calculated
appear in Table 2 below.

Table 2

10

15

20

25

30

COMPOUND	TOXICITY TD ₅₀ μ M
Spermine	2
Spermidine	2
Putrescine	300
(R)-1-aminoindan	175
2709	50
2710	30
2711	60
2715	10
2716	0.7
2720	0.2
2721	1
2722	20
2724	4
2725	20
2728	0.25
2726	1.7
2727	0.4

2. Global Brain Ischemia In Gerbils

Male mongolian gerbils, aged 2.5-5 months, housed 4-8
35 in a cage, were supplied freely with food and water
and maintained at 24°C with a 12 hour day/night cycle.

-53-

For surgery the animals were anesthetized with halothane (1.5% in 100% O₂) and the common carotid arteries exposed bilaterally through a midline ventral neck incision. Each artery was clamped for 5 minutes with aneurysm clips to produce global brain ischemia. The anesthesia was discontinued upon clamping. After 5 minutes the clamps were removed and the wound closed with skin clips.

For treatment, the test compound or vehicle was injected intra-peritoneally (ip) in a volume of 50 microlitres of solvent per 10g body weight. The first injection was given two minutes after clamp removal, and thereafter daily for the next 2 post-operative days (total of 3 injections).

Seven groups each of 4-8 animals were compared:

- I. Control; (sham- or unoperated and saline treated),
- II. Vehicle treated, unoperated (when solvent is other than saline),
- III. Ischemia - untreated or saline treated),
- IV. Ischemia - vehicle treated (when solvent is other than saline),
- V. Ischemia - Test compound treated, one sub-group per concentration tested,
- VI. Ischemia and pentobarbital (40 mg/kg) as positive control).

Analysis of neuronal damage was performed 14 days post-ischemia by counting pyramidal neurons throughout the CA1 layer of the anterior hippocampus in 4 micromolar thick (paraffin) coronal brain sections stained with hematoxylin and eosin.

The Results are shown in the Table 3 below. Percent Protection values represent the fraction of hippocampi protected from ischemia in a group of animals treated with the stated dose of compound.

Table 3

	TEST COMPOUND (n)	DOSE (mg/kg)	% INTACT NEURONS	PERCENT PROTECTION
5	Control Ischemia Pentobarbital (8)		100 0	87.5
	Spermine	10.0		60
10	Spermidine	10.0		42
	Putrescine	10.0		41
15	(R) -1-aminoindan (4) (4) (4)	0.1 1.0 10.0		25.0 25.0 25.0
20	(S) -1-aminoindan (4) (4) (4)	0.1 1.0 10.0		0 50.0 75.0
	2709 (4) (8)	1.0 10.0		12 31
	2710 (4) (8)	1.0 10.0		25 37
25	2711 (4) (12) (5)	1.0 10.0 10.0		50 86 70
	2715 (4) (7)	1.0 10.0		75 50
30	2716 (4) (7)	1.0 10.0		50 0
	2720 (2) (12)	1.0 10.0		- 58
35	2721 (8) (6)	1.0 10.0		12 67
	2722 (4) (4)	1.0 10.0		50 25
	2724 (4) (13)	1.0 10.0		12 56
40	2725 (4) (4)	1.0 10.0		0 25

-55-

5	2728	(5)	1.0		20
		(5)	10.0		80
		(5)	10.0		40
	2726	(4)	10.0		50
	2727	(7)	10.0		28.5

FOCAL CEREBRAL ISCHEMIA IN RATS

10

Ischemia was produced by intraluminal middle cerebral artery (MCA) occlusion in 4-month-old male Wistar rats. The surgery was performed under halothane anesthesia according to Longa et al. (Stroke, 20: 84-91, 1989). Briefly, the left common carotid artery (CCA) was exposed through a ventral neck incision and temporarily clamped, the pterygopalatine artery ligated, the external carotid artery (ECA) cut and a 4-0 nylon monofilament (its tips rounded by heating) was introduced and advanced about 18 mm through the ECA and the internal carotid artery into the anterior cerebral artery to block the origin of the MCA. The monofilament was then secured with a suture around the ECA, the clamp removed from the CCA and the anesthesia discontinued. After 3 h of ischemia without anesthesia, the rats were lightly anesthetized the monofilament withdrawn and re-perfusion commenced. Drugs: 10 mg/kg of 2711 were injected intraperitoneally 5 min. after beginning of re-perfusion and again once daily for the next 2 postoperative days (altogether 3 injections). Saline injected animals served as controls. Body weight was recorded just prior to surgery and then daily. Histology: Animals were killed 3 days after the occlusion

-56-

and the brains were removed and fixed in buffered pH 7.4 formalin. Cross sections (5 μ M thick paraffin sections) were taken at 3 representative levels (4.5mm (A), 6.5mm (B) and 10.5mm (C) anterior to the intraural line) and stained with hematoxylin and eosin. The damaged area, containing damaged pyknotic neurons in the parietal and olfactory cortex and in the striatum, was measured by planimetry under the microscope. Tables 4 and 5 below show the effect of 2711 on the area of brain damage and body weight respectively.

Table 4

15

<u>Brain damage (mm²)</u>			
	A	B	C
Saline	8.7 \pm 2.5	15.2 \pm 3.5	21.0 \pm 4.7
TV-2711	1.4 \pm 0.5	4.6 \pm 3.3	7.4 \pm 3.8

30

Table 5


5	Body Weight					
	Saline (%)	(6)	425±5 (100)	377±5 (89)	357±6 (84)	342±6 (81)
10	2711 (%)	(5)	430±3 (100)	396±6 (92)	384±5 (89)	379±6 (88)

The effect of 2711 on the outcome of focal
15 cerebral ischemia in rats in terms of damage
to the brain (table 1) and effect on body
weight (table 2). Results are expressed in
mean±S.E.M. values.

-58-

What is claimed is:

1. A compound of the formula:



wherein n is 1 or 2;

R_1 and R_2 are each independently hydrogen, hydroxy, substituted or unsubstituted C_{1-4} alkyl, substituted or unsubstituted C_{1-4} alkoxy or halogeno;

15 R_5 is hydrogen, substituted or unsubstituted C_{1-4} alkyl, amino or NR_6R_7 ;

R_3 , R_4 , R_6 and R_7 are each independently hydrogen or $(CH_2)_m - [NH(CH_2)_x]_y - NHR_{11}$ wherein m is from 0-5, y is from 0-5 and each x is independently from 1-5 and R_{11} is hydrogen or $C(NH)NH_2$;

20 provided that when R_3 and R_4 are both hydrogen, R_5 is amino or NR_6R_7 ; pharmaceutically acceptable salts and optically active isomers thereof.

25 2. The compound of claim 1, wherein n is 1.

3. The compound of claim 1, wherein R_1 , R_2 and R_5 are hydrogen.

30 4. The compound of claim 1, wherein R_3 is $(CH_2)_m - [NH(CH_2)_x]_y - NHR_{11}$ and R_4 is hydrogen.

5. The compound of claim 4, wherein R_{11} is hydrogen.

35 6. The compound of claim 4, wherein m is 3 or 4.

7. The compound of claim 4, wherein y is 0, 1 or 2.

-59-

8. The compound of claim 4, wherein each x is 3 or 4.
9. The compound of claim 4, wherein y is greater than 1, and each x is 3 or 4.
- 5 10. The compound of claim 4, wherein R_{11} is $C(NH)NH_2$.
11. The compound of claim 1, wherein both R_3 and R_4 are $(CH_2)_m - [NH(CH_2)_x]_y - NHR_{11}$.
- 10 12. The compound of claim 11, wherein R_{11} is hydrogen.
13. The compound of claim 11, wherein m is 3 or 4.
- 15 14. The compound of claim 11, wherein y is 0, 1 or 2.
15. The compound of claim 14, wherein y is 0.
16. The compound of claim 11, wherein R_{11} is $C(NH)NH_2$.
- 20 17. The compound of claim 1, wherein R_3 and R_4 are hydrogen; and R_5 is amino or NR_6R_7 .
18. The compound of claim 17, wherein m is 3 or 4; y is 0, 1 or 2; and each x is 3 or 4.
- 25 19. The compound of claim 17, wherein R_5 is amino.
20. The compound of claim 17, wherein R_5 is NR_6R_7 .
- 30 21. The compound of claim 20, wherein R_6 is hydrogen; and R_7 is $(CH_2)_m - [NH(CH_2)_x]_y - NHR_{11}$.
22. The compound of claim 21, wherein m is 3 or 4; y is 0; and R_{11} is hydrogen.
- 35 23. The compound of claim 1, wherein R_5 is trans with

-60-

respect to NR_3R_4 .

24. The compound of claim 1, wherein R_5 is cis with respect to NR_3R_4 .
- 5
25. The compound of claim 1, wherein the ring carbon to which NR_3R_4 is bound is in the R configuration.
26. The compound of claim 1, wherein the ring carbon to which NR_3R_4 is bound is in the S configuration.
- 10
27. The compound of claim 1, selected from the group consisting of:
- 15
- N-(3-aminopropyl)-1-aminoindan;
- N-(4-aminopropyl)-1-aminoindan;
- N,N-di(4-amino-n-butyl)-1-aminoindan;
- (R)-N,N-di(4-amino-n-butyl)-1-aminoindan;
- (S)-N,N-di(4-amino-n-butyl)-1-aminoindan;
- 20
- N-(4-amino-n-butyl)-N-(3-amino-n-propyl)-1-aminoindan;
- N,N-di(4-amino-n-butyl)-1-amino-1,2,3,4-tetrahydronaphthalene;
- N-[N-(4-aminobutyl)-3-aminopropyl]-1-aminoindan;
- N-[N-(3-aminopropyl)-4-aminobutyl]-1-aminoindan;
- 25
- N-{[N-(3-aminopropyl)-4-aminobutyl]}-N'-3-aminopropyl-1-aminoindan;
- N-{[N-(4-aminobutyl)-3-aminopropyl]}-N'-4-aminobutyl-1-aminoindan;
- cis N,N'-diacetyl-1,3-diaminoindan;
- 30
- trans N,N'-diacetyl-1,3-diaminoindan;
- cis 1,3-diaminoindan;
- trans 1,3-diaminoindan;
- N-(4-aminobutyl)-1,3-diaminoindan;
- 35
- N,N'-di(4-aminobutyl)-1,3-diaminoindan;
- N-(3-aminopropyl)-N-[N-(3-aminopropyl)-4-aminobutyl]-1-aminoindan;

-61-

N,N-di(4-guanidinobutyl)-1-aminoindan;
N-(4-guanidinobutyl)-1-aminoindan,
or a pharmaceutically acceptable salt thereof.

- 5 28. A pharmaceutical composition comprising a
therapeutically effective amount of the compound of
claim 1 and a pharmaceutically acceptable carrier.
- 10 29. The pharmaceutical composition of claim 28, wherein
the pharmaceutically acceptable carrier is a solid
and the pharmaceutical composition is a tablet.
- 15 30. The pharmaceutical composition of claim 28, wherein
the therapeutically effective amount is from about
1 mg to about 1000 mg.
- 20 31. The pharmaceutical composition of claim 30, wherein
the therapeutically effective amount is from about
10 mg to about 100 mg.
- 25 32. The pharmaceutical composition of claim 28, wherein
the pharmaceutically acceptable carrier is a liquid
and the pharmaceutical composition is an injectable
solution.
- 30 33. The pharmaceutical composition of claim 32, wherein
the therapeutically effective amount is from about
1 mg/ml to about 1000 mg/ml.
- 35 34. The pharmaceutical composition of claim 33, wherein
the therapeutically effective amount is from about
10 mg/ml to about 100 mg/ml.
- 35 35. The pharmaceutical composition of claim 28, wherein
the carrier is a gel and the pharmaceutical
composition is a suppository.

-62-

36. A method of treating acute neurological traumatic disorder or neurotrauma in a subject comprising administering to the subject a therapeutically effective amount of the compound of claim 1 or pharmaceutically acceptable salt thereof, effective to treat the acute neurological traumatic disorder or neurotrauma in the subject.
37. A method of treating a subject afflicted with a neurodegenerative disease which comprises administering to the subject an amount of the compound of claim 1 or pharmaceutically acceptable salt thereof, effective to treat the neurodegenerative disease in the subject.
38. A method of treating a subject afflicted with a neurotoxic injury which comprises administering to the subject an amount of the compound of claim 1 or pharmaceutically acceptable salt thereof, effective to treat the neurotoxic injury in the subject.
39. A method of treating a subject afflicted with brain ischemia which comprises administering to the subject an amount of the compound of claim 1 or pharmaceutically acceptable salt thereof, effective to treat brain ischemia in the subject.
40. A method of treating a subject afflicted with a head trauma injury which comprises administering to the subject an amount of the compound of claim 1 or pharmaceutically acceptable salt thereof, effective to treat the head trauma injury in the subject.
41. A method of treating a subject afflicted with a spinal trauma injury which comprises administering to the subject an amount of the compound of claim 1 or pharmaceutically acceptable salt thereof,

-63-


effective to treat the spinal trauma injury in the subject.

- 5 42. A method of preventing nerve damage in a subject which comprises administering to the subject an amount of the compound of claim 1 or pharmaceutically acceptable salt thereof, effective to prevent nerve damage in the subject.

1/5

FIGURE 1A


Scheme I



2/5

FIGURE 1B


Scheme I cont.



3/5


FIGURE 2

Scheme II




4/5

FIGURE 3



5/5

FIGURE 4



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/11318

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/15, 31/155; C07C 243/20, 247/14, 279/12, 281/16.

US CL : 514/616, 632, 634, 657; 564/222, 228, 236, 238, 428.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/616, 632, 634, 657; 564/222, 228, 236, 238, 428.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE - structure search.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3,953,506 A (SPICER ET AL.) 27 April 1976 (27.04.76), column 27, lines 30-35.	1-3, 17, 19, 20, 23-26, 28-35
X	US 4,792,628 A (OSHIRO ET AL.) 20 December 1988 (20.12.88), column 30, example 20.	1-3, 17, 19, 20, 23-26, 28-35
X	HUEBNER et al. A Product of the Chemical Reduction of <i>o</i> -Benzenediacetonitrile. Journal of Organic Chemistry. December 1962, Vol. 27, pages 4465-4469, especially compounds VIII and IXa and the examples on pages 4468 and 4469.	1-5, 7, 23-26

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

02 AUGUST 1996

Date of mailing of the international search report

21 AUG 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RICHARD L. RAYMOND

Telephone No. (703) 308-1235